ELSEVIER

Neuroscience Letters 350 (2003) 195–198

Neuroscience Letters

www.elsevier.com/locate/neulet

Author Index to Volume 350

Altura, B.M., see Li, J., 173
Altura, B.T., see Li, J., 173
Amano, M., Iigo, M., Ikuta, K., Kitamura, S., Yamamori, K., Daily variations in melatonin binding sites in the masu salmon brain, 9
Amiridis, I.G., Hatzitaki, V., Arabatzi, F., Age-induced modifications of static postural control in humans, 137
Amouyel, P., see Helbecque, N., 181
Aoki, T., Narita, M., Ohnishi, O., Mizuo, K., Narita, M., Yajima, Y., Suzuki, T., Disruption of the type 1 inositol 1,4,5-trisphosphate receptor gene suppresses the morphine-induced antinociception in the mouse, 69
Arabatzi, F., see Amiridis, I.G., 137
Archetti, S., see Poli, M., 77
Auindx for Nsl 350/1
Ayabe, M., see Kinoshita, T., 169

Bach, A., see Kuner, R., 153
Bachelin, C., see Pagany, M., 165
Bae, S.-r., see Shin, D.H., 149
Bagdy, G., see Juhasz, G., 56
Bagust, J., see Maile, R., 190
Baier, M., see Schwarz, A., 187
Bamme, T., see Schwarz, A., 187
Barish, M.E., see Ring, R.H., 73
Baron van Evercooren, A., see Pagany, M., 165
Beach, T.G., see Newell, A.J., 178
Bellido, I., Gomez-Luque, A., Garcia-Carrera, P., Rius, F., de la Cuesta, F.S., Female rats show an increased sensibility to the forced swim test depressive-like stimulus in the hippocampus and frontal cortex 5-HT_{1A} receptors, 145
Biello, S.M., see Dafters, R.I., 117
Buchner, H., see Gobbelé, R., 97
Burwinkel, M., see Schwarz, A., 187

Cairns, N.J., see Culpan, D., 61
Camboni, D., Roskoden, T., Schwegler, H., Effect of early thyroxine treatment on brain-derived neurotrophic factor mRNA expression and protein amount in the rat medial septum/diagonal band of Broca, 141
Castrén, E., see Väisänen, J., 1
Catania, S., Germanà, A., Laurà, R., Gonzalez-Martinez, T., Ciriaco, E., Vega, J.A., The crypt neurons in the olfactory epithelium of the adult zebrafish express TrkA-like immunoreactivity, 5
Cha, C.I., see Shin, D.H., 149
Cheema, S.S., see Turner, B.J., 132
Cheung, J., see Yang, J.D., 25
Chung, Y.H., see Shin, D.H., 149
Ciriaco, E., see Catania, S., 5
Cocozza, S., see Turano, M., 184
Contreras, R.J., see Curtis, K.S., 46
Cottel, D., see Helbecque, N., 181
Culpan, D., MacGowan, S.H., Ford, J.M., Nicoll, J.A.R., Griffin, W.S., Dewar, D., Cairns, N.J., Hughes, A., Kehoe, P.G., Wilcock, G.K., Tumour necrosis factor-α gene polymorphisms and Alzheimer's disease, 61
Curio, G., see Gobbelé, R., 97
Curtis, K.S., Krause, E.G., Contreras, R.J., Cardiovascular function and circadian patterns in rats after area postrema lesions or prolonged food restriction, 46

Dafters, R.I., Biello, S.M., The effect of 3,4-methylenedioxymethamphetamine ('Ecstasy') on serotonergic regulation of the mammalian circadian clock mechanism in rats: the role of dopamine and hyperthermia, 117
Darrington, R.S., Leigh, P.N., Gallo, J.-M., Protective effects of estrogens on polyglutamine-expanded androgen receptor aggregation in mice, 37
Darvas, F., see Gobbelé, R., 97
Daum, I., see Suchan, B., 101
De Biase, I., see Turano, M., 184
de la Cuesta, F.S., see Bellido, I., 145
Dewar, D., see Culpan, D., 61
Domijan, D., A mathematical model of persistent neural activity in human prefrontal cortex for visual feature binding, 89
Dreyer, J., Hirlinger, D., Müller-Esterl, W., Oess, S., Kuner, R., Spinal upregulation of the nitric oxide synthase-interacting protein NOSIP in a rat model of inflammatory pain, 13

Endo, S., Launey, T., Nitric oxide activates extracellular signal-regulated kinase 1/2 and enhances declustering of ionotropic glutamate receptor subunit 2/3 in rat cerebellar Purkinje cells, 122

Faludi, G., see Juhasz, G., 56
Feng, G.Y., see Yang, J.D., 25
Ferger, B., see Kuner, R., 153
Finazzi, D., see Poli, M., 77
Ford, J.M., see Culpan, D., 61
Fujimoto, H., see Kinoshita, T., 169

Gallo, J.-M., see Darrington, R.S., 37
Gao, C., see Ring, R.H., 73
Garcia-Carrera, P., see Bellido, I., 145
Gatta, L.B., see Poli, M., 77
Géranton, S.M., Heal, D.J., Stanford, S.C., Noradrenaline clearance differs in the frontal cortex and hypothalamus: a dual-probe microdialysis study in freely-moving rats, 157
Germanà, A., see Catania, S., 5
Goadsby, P.J., see Knight, Y.E., 85
Gobbelé, R., Waberski, T.D., Thyerlei, D., Thissen, M., Darvas, F., Klostermann, F., Curio, G., Buchner, H., Functional dissociation of a subcortical and cortical component of high-frequency oscillations in human somatosensory evoked potentials by motor interference, 97

doi:10.1016/S0304-3940(03)01053-X

Gomez-Luque, A., see Bellido, I., 145
Gonzalez-Martinez, T., see Catania, S., 5
Griffin, W.S., see Culpan, D., 61

Han, J.-S., see Huang, C., 93
Hatzitaki, V., see Amiridis, I.G., 137
He, L., see Yang, J.D., 25
Heal, D.J., see Géranton, S.M., 157
Helbecque, N., Hermant, X., Cottel, D., Amouyel, P., The role of matrix metalloproteinase-9 in dementia, 181
Henklein, P., see Schwarz, A., 187
Hermant, X., see Helbecque, N., 181
Hirlinger, D., see Dreyer, J., 13
Huang, C., Long, H., Shi, Y.-S., Han, J.-S., Wan, Y., Nocistatin potentiates electroacupuncture antinociceptive effects and reverses chronic tolerance to electroacupuncture in mice, 93
Huang, N.-K., Adenosine A_{2A} receptors regulate oxidative stress formation in rat pheochromocytoma PC12 cells during serum deprivation, 127
Hughes, A., see Culpan, D., 61

Iigo, M., see Amano, M., 9
Ikeda, S.-i., see Ishii, K., 161
Ikuta, K., see Amano, M., 9
Imaizumi, T., see Kinoshita, T., 169
Inagaki, C., see Zhang, N.-Y., 105
Inui, T., see Okada, T., 21
Ishibashi, M., see Yasui, F., 66
Ishii, K., Tokuda, T., Matsushima, T., Miya, F., Shoji, S., Ikeda, S.-i., Tamaoka, A., Pravastatin at 10 mg/day does not decrease plasma levels of either amyloid-β (Aβ) 40 or Aβ 42 in humans, 161
Isobe, K., see Nanmoku, T., 33

Jagodic, M., see Pagany, M., 165
Jakus, R., see Juhasz, G., 56
Joo, K.M., see Shin, D.H., 149
Juhasz, G., Zsombok, T., Laszik, A., Jakus, R., Faludi, G., Sotonyi, P., Bagdy, G., Despite the general correlation of the serotonin transporter gene regulatory region polymorphism (*5-HTTLPR*) and platelet serotonin concentration, lower platelet serotonin concentration in migraine patients is independent of the *5-HTTLPR* variants, 56

Kalitzin, S., Suffczynski, P., Comments on "Correlations between brain electrical activities of two spatially separated human subjects", Wackermann et al., Neurosci. Lett. 336 (2003) 60–64, 193
Kanemitsu, H., Tomiyama, T., Mori, H., Human neprilysin is capable of degrading amyloid β peptide not only in the monomeric form but also the pathological oligomeric form, 113
Kasai, M., see Sugiura, T., 109
Kato, S., see Yamashita, S., 17
Katsuya, H., see Sugiura, T., 109
Kawakami, Y., see Nanmoku, T., 33
Kehoe, P.G., see Culpan, D., 61
Kenchappa, R.S., Ravindranath, V., γ-Glutamyl cysteine synthetase is up-regulated during recovery of brain mitochondrial complex I following neurotoxic insult in mice, 51
Kim, J., see Shin, D.H., 149
Kinoshita, T., Imaizumi, T., Miura, Y., Fujimoto, H., Ayabe, M., Shoji, H., Okamoto, Y., Takashima, H., Osame, M., Nakagawa, M., A case of adult-onset Alexander disease with Arg416Trp human glial fibrillary acidic protein gene mutation, 169
Kitagawa, K., see Zhang, N.-Y., 105
Kitamura, S., see Amano, M., 9
Klostermann, F., see Gobbelé, R., 97
Knight, Y.E., Levy, M.J., O'Shaughnessy, C.T., Goadsby, P.J., Prostaglandin E2 injected into the posterior hypothalamus has no effect on trigeminal nociception in the rat, 85
Kojo, S., see Yasui, F., 66
Konishi, J., see Okada, T., 21
Koponen, E., see Väisänen, J., 1
Krätke, O., see Schwarz, A., 187
Krause, E.G., see Curtis, K.S., 46
Kuner, R., see Dreyer, J., 13
Kuner, R., Teismann, P., Trutzel, A., Naim, J., Richter, A., Schmidt, N., von Ahsen, O., Bach, A., Ferger, B., Schneider, A., TorsinA protects against oxidative stress in COS-1 and PC12 cells, 153

Lachapelle, F., see Pagany, M., 165
Laszik, A., see Juhasz, G., 56
Launey, T., see Endo, S., 122
Laurà, R., see Catania, S., 5
Lee, E., see Shin, D.H., 149
Leigh, P.N., see Darrington, R.S., 37
Levy, M.J., see Knight, Y.E., 85
Li, J., Su, J., Li, W., Liu, W., Altura, B.T., Altura, B.M., Peroxynitrite induces apoptosis in canine cerebral vascular muscle cells: possible relation to neurodegenerative diseases and strokes, 173
Li, W., see Li, J., 173
Linington, C., see Pagany, M., 165
Liu, W., see Li, J., 173
Liu, Z.-W., see Yang, S., 81
Lo Casale, M.S., see Turano, M., 184
Long, H., see Huang, C., 93
Lopes, E.C., see Turner, B.J., 132

MacGowan, S.H., see Culpan, D., 61
Maile, R., Walker, R.J., Sharma, R.P., Bagust, J., Effects of nociceptin and analogues of nociceptin upon spontaneous dorsal root activity recorded from an in vitro preparation of rat spinal cord, 190
Makita, K., see Takei, T., 41
Matsugo, S., see Yasui, F., 66
Matsushima, T., see Ishii, K., 161
Mita, S., see Yamashita, S., 17
Miura, Y., see Kinoshita, T., 169
Miya, F., see Ishii, K., 161
Mizumura, K., see Sugiura, T., 109
Mizuo, K., see Aoki, T., 69
Monticelli, A., see Turano, M., 184
Mori, H., see Kanemitsu, H., 113
Müller-Esterl, W., see Dreyer, J., 13
Murakoshi, T., see Takei, T., 41
Mutoh, S., see Ochi, T., 29

Naim, J., see Kuner, R., 153
Nakagawa, M., see Kinoshita, T., 169
Nakai, T., see Nanmoku, T., 33
Nakai, T., see Okada, T., 21
Nanmoku, T., Takekoshi, K., Isobe, K., Kawakami, Y., Nakai, T., Okuda, Y., Prolactin-releasing peptide stimulates catecholamine release but not proliferation in rat pheochromocytoma PC12 cells, 33
Narita, M., see Aoki, T., 69

Narita, M., see Aoki, T., 69
Newell, A.J., Sue, L.I., Scott, S., Rauschkolb, P.K., Walker, D.G., Potter, P.E., Beach, T.G., Thiorphan-induced neprilysin inhibition raises amyloid β levels in rabbit cortex and cerebrospinal fluid, 178
Nicoll, J.A.R., see Culpan, D., 61
Nishizawa, S., see Okada, T., 21

Ochi, T., Ohkubo, Y., Mutoh, S., Effect of systemic FR140423, a new analgesic compound, in a rat model of postoperative pain: contribution of delta-opioid receptors, 29
Oess, S., see Dreyer, J., 13
Ohama, E., see Yamashita, S., 17
Ohkubo, Y., see Ochi, T., 29
Ohnishi, O., see Aoki, T., 69
Okada, T., Tanaka, S., Nakai, T., Nishizawa, S., Inui, T., Yonekura, Y., Konishi, J., Sadato, N., Facial recognition reactivates the primary visual cortex: an functional magnetic resonance imaging study in humans, 21
Okado, H., see Yamashita, S., 17
Okamoto, Y., see Kinoshita, T., 169
Okuda, Y., see Nanmoku, T., 33
Olsson, T., see Pagany, M., 165
Oomura, Y., see Yasui, F., 66
Osame, M., see Kinoshita, T., 169
O'Shaughnessy, C.T., see Knight, Y.E., 85

Padovani, A., see Poli, M., 77
Pagany, M., Jagodic, M., Schubart, A., Pham-Dinh, D., Bachelin, C., Baron van Evercooren, A., Lachapelle, F., Olsson, T., Linington, C., Myelin oligodendrocyte glycoprotein is expressed in the peripheral nervous system of rodents and primates, 165
Pham-Dinh, D., see Pagany, M., 165
Pianese, L., see Turano, M., 184
Poli, M., Gatta, L.B., Archetti, S., Padovani, A., Albertini, A., Finazzi, D., Association analysis between anterior-pharynx defective-1 genes polymorphisms and Alzheimer's disease, 77
Potter, P.E., see Newell, A.J., 178

Rauschkolb, P.K., see Newell, A.J., 178
Ravindranath, V., see Kenchappa, R.S., 51
Richter, A., see Kuner, R., 153
Riemer, C., see Schwarz, A., 187
Ring, R.H., Valo, Z., Gao, C., Barish, M.E., Singer-Sam, J., The *Cdkn1a* gene ($p21^{Waf1/Cip1}$) is an inflammatory response gene in the mouse central nervous system, 73
Rius, F., see Bellido, I., 145
Roskoden, T., see Camboni, D., 141
Ruggiero, G., see Turano, M., 184

Saarelainen, T., see Väisänen, J., 1
Sadato, N., see Okada, T., 21
Saegusa, H., see Takei, T., 41
Sasaki, K., see Yasui, F., 66
Schmidt, N., see Kuner, R., 153
Schneider, A., see Kuner, R., 153
Schubart, A., see Pagany, M., 165
Schultz, J., see Schwarz, A., 187
Schwarz, A., Krätke, O., Burwinkel, M., Riemer, C., Schultz, J., Henklein, P., Bamme, T., Baier, M., Immunisation with a synthetic prion protein-derived peptide prolongs survival times of mice orally exposed to the scrapie agent, 187
Schwegler, H., see Camboni, D., 141
Scott, S., see Newell, A.J., 178
Sharma, R.P., see Maile, R., 190
Shi, Y.-S., see Huang, C., 93
Shin, D.H., Lee, E., Joo, K.M., Kim, J., Bae, S.-r., Chung, Y.H., Cha, C.I., Immunocytochemical study on the distribution of c-myb in the central nervous system of the transgenic mice expressing a human copper/zinc superoxide dismutase mutation, 149
Shoji, H., see Kinoshita, T., 169
Shoji, S., see Ishii, K., 161
Singer-Sam, J., see Ring, R.H., 73
Sotonyi, P., see Juhasz, G., 56
St. Clair, D., see Yang, J.D., 25
Stanford, S.C., see Géranton, S.M., 157
Su, J., see Li, J., 173
Suchan, B., Zoppelt, D., Daum, I., Frontocentral negativity in electroencephalogram reflects motor response evaluation in humans on correct trials, 101
Sue, L.I., see Newell, A.J., 178
Suffczynski, P., see Kalitzin, S., 193
Sugiura, T., Kasai, M., Katsuya, H., Mizumura, K., Thermal properties of acid-induced depolarization in cultured rat small primary afferent neurons, 109
Suzuki, T., see Aoki, T., 69

Takashima, H., see Kinoshita, T., 169
Takei, T., Saegusa, H., Zong, S., Murakoshi, T., Makita, K., Tanabe, T., Increased sensitivity to halothane but decreased sensitivity to propofol in mice lacking the N-type Ca^{2+} channel, 41
Takekoshi, K., see Nanmoku, T., 33
Tamaoka, A., see Ishii, K., 161
Tammaro, A., see Turano, M., 184
Tanabe, T., see Takei, T., 41
Tanaka, S., see Okada, T., 21
Teismann, P., see Kuner, R., 153
Thissen, M., see Gobbelé, R., 97
Thyerlei, D., see Gobbelé, R., 97
Tokuda, T., see Ishii, K., 161
Tomiyama, T., see Kanemitsu, H., 113
Trutzel, A., see Kuner, R., 153
Turano, M., Tammaro, A., De Biase, I., Lo Casale, M.S., Ruggiero, G., Monticelli, A., Cocozza, S., Pianese, L., 3-Nitropropionic acid increases frataxin expression in human lymphoblasts and in transgenic rat PC12 cells, 184
Turner, B.J., Lopes, E.C., Cheema, S.S., Neuromuscular accumulation of mutant superoxide dismutase 1 aggregates in a transgenic mouse model of familial amyotrophic lateral sclerosis, 132

Uchino, M., see Yamashita, S., 17

Väisänen, J., Saarelainen, T., Koponen, E., Castrén, E., Altered trkB neurotrophin receptor activation does not influence the *N*-methyl-D-aspartate receptor antagonist-mediated neurotoxicity in mouse posterior cingulate cortex, 1
Valo, Z., see Ring, R.H., 73
Vega, J.A., see Catania, S., 5
von Ahsen, O., see Kuner, R., 153

Waberski, T.D., see Gobbelé, R., 97
Wackermann, J., Correlations between brain electrical activities of two

spatially separated human subjects. Reply to the commentary by S. Kalitzin and P. Suffczynski, 194
Walker, D.G., see Newell, A.J., 178
Walker, R.J., see Maile, R., 190
Wan, Y., see Huang, C., 93
Wilcock, G.K., see Culpan, D., 61
Wu, B., see Zhang, N.-Y., 105

Yajima, Y., see Aoki, T., 69
Yamamori, K., see Amano, M., 9
Yamashita, S., Mita, S., Kato, S., Okado, H., Ohama, E., Uchino, M., Bcl-2 expression using retrograde transport of adenoviral vectors inhibits cytochrome c-release and caspase-1 activation in motor neurons of mutant superoxide dismutase 1 (G93A) transgenic mice, 17
Yang, J.D., Feng, G.Y., Zhang, J., Cheung, J., St. Clair, D., He, L., Apolipoprotein E −491 promoter polymorphism is an independent risk factor for Alzheimer's disease in the Chinese population, 25
Yang, S., Liu, Z.-W., Zhou, W.-X., Zhang, Y.-X., Thymosin alpha-1 modulate excitatory synaptic transmission in cultured hippocampal neurons in rats, 81
Yasui, F., Ishibashi, M., Matsugo, S., Kojo, S., Oomura, Y., Sasaki, K., Brain lipid hydroperoxide level increases in senescence-accelerated mice at an early age, 66
Yonekura, Y., see Okada, T., 21

Zhang, J., see Yang, J.D., 25
Zhang, N.-Y., Kitagawa, K., Wu, B., Inagaki, C., Soybean-derived phosphatidylinositol recovers amyloid β protein-induced neurotoxicity in cultured rat hippocampal neurons, 105
Zhang, Y.-X., see Yang, S., 81
Zhou, W.-X., see Yang, S., 81
Zong, S., see Takei, T., 41
Zoppelt, D., see Suchan, B., 101
Zsombok, T., see Juhasz, G., 56